



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**2825. Proposed by the late L. G. WELD.**

A ball, having a coefficient of resilience  $\alpha$ , strikes a rigid plane surface, inclined at an angle  $\theta$  from the horizontal, after falling through a height  $h$ . What is the distance from the first to the second point of impact with the plane?

**2826. Proposed by ALBERT A. BENNETT, University of Texas.**

As a standard form for a square non-singular symmetric matrix under certain transformations, may be taken the form in which only the elements in the secondary diagonal are different from zero, and each of these is equal to unity. Analogously, as a standard form for a square non-singular skew-symmetric matrix (and hence incidentally of even order), may be taken the form in which only the elements of the secondary diagonal are different from zero, while the half of these which are towards the upper right-hand corner are each minus one, and the remaining half towards the lower left-hand corner, are each plus one. Denote both of these standard matrices by  $N$ .

Give simple parallel proofs that if  $M$  be given as non-singular and symmetric or non-symmetric as the case may be, a matrix  $P$  exists such that, with the usual notation

$$M = PNP'.$$

**2827. Proposed by B. F. FINKEL, Drury College.**

Find the equation of the envelope of the system of circles inscribed in a triangle having a given base and a given altitude.

**2828. Proposed by T. M. BLAKSLEE, Ames, Iowa.**

On page 72 of R. B. Hayward's *The Algebra of Coplanar Vectors and Trigonometry* occurs the sentence: "It will be a good exercise for the student to show that  $\cos(90^\circ/7) = \frac{1}{2}\sqrt{x_1}$ , where  $x_1$  is the greatest root of the equation,

$$x^3 - 7x^2 + 14x - 7 = 0."$$

(1) Do not merely verify but deduce the equation and find  $x_1$ . (2) Deduce the  $x$ -equation ( $x_1, x_2, x_3, x_4$ , the roots) such that its greatest root  $x_1$  gives  $\cos(90^\circ/9) = \cos 10^\circ = \frac{1}{2}\sqrt{x_1}$ . (3) Of what angles are  $\frac{1}{2}\sqrt{x_1}, \dots, \frac{1}{2}\sqrt{x_4}$ , in (2), the cosines? Develop a method of writing out at once  $\cos(nv)$  in terms of powers of  $\cos v$  if these are given for  $(n-1)v$  and  $(n-2)v$ . The same for  $\sin(nv)$ . (4) Use the results of (2) and (3) to find the number of degrees in a radian. Hence, find  $\pi$  from radian instead of radian from  $\pi$  as is usual.

**SOLUTIONS OF PROBLEMS.****411 (Algebra) [1914, 121; 1919, 268, 459]. Proposed by V. M. SPUNAR, Chicago, Ill.**

Determine  $x_1, x_2, x_3 \dots x_p$  from the equations:

$$\begin{array}{ccccccc} x_1 + x_2 & + x_3 & + \dots + x_p & = a_0, \\ b_1 x_1 + b_2 x_2 & + b_3 x_3 & + \dots + b_p x_p & = a_1, \\ b_1^2 x_1 + b_2^2 x_2 & + b_3^2 x_3 & + \dots + b_p^2 x_p & = a_2, \\ \dots & \dots & \dots & \dots \\ b_1^{p-1} x_1 + b_2^{p-1} x_2 & + b_3^{p-1} x_3 & + \dots + b_p^{p-1} x_p & = a_{p-1}. \end{array}$$

A solution has been sent in by P. J. DA CUNHA, University of Lisbon, Portugal, in which the analysis is very much the same as that previously printed, but he also considers the case when some of the  $b$ 's are equal. For example,

if  $p = 3$  and  $b_1 = b_2 \geq b_3$ , we must have

$$\begin{vmatrix} 1 & 1 & a_0 \\ b_1 & b_3 & a_1 \\ b_1^2 & b_3^2 & a_2 \end{vmatrix} = 0.$$

If this condition is satisfied, the expressions for  $x_1$  and  $x_2$  in the general solution take indeterminate forms, and there will be an infinite number of solutions given by

$$x_1 + x_2 = \frac{a_0 b_3 - a_1}{b_3 - b_1}, \quad x_3 = \frac{a_0 b_1 - a_1}{b_1 - b_3}.$$